ALL-RISK TRIAGE TAGS AND RIBBONS





GOAL:


To establish rapid, uniform, and common means of triage identification for all patients involved in a multi-casualty incident in EDC.



All patients involved in an MCI will be triaged using the START or JUMPSTART evaluation method. Patients should FIRST be triaged using the appropriate colored ribbon, a triage tag will then be applied later during the treatment



- > START TRIAGE:
- Respirations: >30
- Perfusion: Cap refill >2 sec.
- Mentation: Cannot follow simple commands
- ► JUMPSTART TRIAGE: (pediatric):
- Respirations: < 15 or >45 (5 mouth to mask ventilations if apneic and pulse present)
- Perfusion: No palpable pulse
- ▶ Mentation: Posturing/unresponsive



Triage ribbon is dispensed from a fanny bag dispenser that will be carried on all first out emergency response vehicles in EDC. This step replaces the initial use of triage tags and will save considerable time in the initial triage of victims during a multi-casualty incident.



The first responder who initially triages a victim will be responsible for securely fastening a 2 foot piece of triage ribbon on the patient's wrist or ankle with the appropriate color (black, red, yellow, or green)



Each time a patient is triaged you must mark the Triage Count Card located in the top of the fanny bag.





The completed Triage Count Cards are then given to the Triage Unit Leader where they are counted and the totals are communicated to the IC/MGS.



TriageTags.com



As patients are moved into treatment areas they can be re-triaged and have triage tags applied.



Triage tags are stored in the DMS MCI position

bags:



HOW TO USE TRIAGE TAGS:

They can be worn as a necklace or

tied to an wrist or ankle.



If the patient is contaminated with a hazardous material or substance you must first initiate decontamination measures, then the patient may be triaged. Tear the "contaminated" tab off and place it with the patients clothing and other contaminated items. You can then perform START triage and determine which category to place the patient in.



► Using the TAGS:

For MINOR patients the tag should look like this:



Using the TAGS:

For DELAYED patients the tag should look like this:




► Using the TAGS:

For IMMEDIATE patients the tag should look like this:



The separated tags are placed in the respective Triage Tag Receipt Holders by the Treatment Area Managers.




Using the TAGS:

For DECEASED patients the tag should look like this:



The wristband can be placed on the patient or just left on the triage tag for the hospital to use later.



The Ambulance Receipt should be collected by the Air/Ground Ambulance Coordinator and be placed in the Transportation Sheet Holder form.



The Personal Property Receipt can be placed with the patient's clothing or other belongings.

